



CATARACT EYE DROP CHART



Tobramycin



Prolensa
 Bromsite
 Ketorolac

- Don't forget to pick up your prescription order at your pharmacy at least one week before your surgery.
- Helpful Hint: Using drops at meal times will help you remember.
- If we have given you coupons for your eye drops, please be sure to take them with you to the pharmacy.
- Your prescriptions will be sent to the pharmacy electronically unless otherwise noted.
- When applying drops wait five minutes between each drop.

Your OPERATIVE EYE is your **RIGHT EYE** or **LEFT EYE**

MEDICATION INSTRUCTIONS:

Start using the drops with the TAN/OFF WHITE CAP and GRAY CAP on _____.
(2 Days Prior to surgery)

Use the chart to track drops by filling in a circle every time you use a medication drop.

BEFORE SURGERY		
DROPS	DAY 1	DAY 2
Tobramycin	<input type="radio"/> <input type="radio"/> <input type="radio"/> <input type="radio"/>	<input type="radio"/> <input type="radio"/> <input type="radio"/> <input type="radio"/>
Prolensa/Bromsite or Ketorolac	<input type="radio"/>	<input type="radio"/>
	<input type="radio"/> <input type="radio"/> <input type="radio"/> <input type="radio"/>	<input type="radio"/> <input type="radio"/> <input type="radio"/> <input type="radio"/>

Please see the back of this page for after surgery instructions